

Literature cited in Table 21.1

- Agostini I, Holzmann I, Di Biteti MS (2010) Are howler monkey species ecologically equivalent? Trophic niche overlap in syntopic *Alouatta guariba clamitans* and *Alouatta caraya*. Am J Primatol 72:173–186
- Aguiar LM, Reis NR, Ludwig G, Rocha VJ (2003) Dieta, área de vida, vocalizações e estimativas populacionais de *Alouatta guariba* em um remanescente florestal no norte do estado do Paraná. Neotrop Primates 11:78–86
- Aldana-Saavedra JP (2009) Feeding ecology and seed dispersal by *Ateles hybridus*, *Alouatta seniculus* and *Cebus albifrons* in a fragmented area at San Juan del Carare, Colombia. MSc thesis, International Master Programme at the Swedish Biodiversity Centre, Uppsala
- Almeida-Silva B, Guedes PG, Boubli JP, Strier KB (2005) Deslocamento terrestre e o comportamento de beber em um grupo de barbados (*Alouatta guariba clamitans* Cabrera, 1940) em Minas Gerais, Brasil. Neotrop Primates 13:1–3
- Alves IMSC, Guix JC (1992) Feeding habits of *Alouatta caraya* in a semi-natural Area (SE Brazil). Mammalia 56:469–472
- Alves SL, Zaú AS (2007) Aspectos ecológicos de *Alouatta guariba clamitans* Cabrera, 1940 na área de relevante interesse ecológico Floresta da Cicutá, Rio de Janeiro, Brasil. Neotrop Primates 14:127–130
- Arditi SI (1992) Variaciones estacionales en la actividad y dieta de *Aotus azarae* y *Alouatta caraya* en Formosa, Argentina. Boletín Primatológico Latinoamericano 3:11–30
- Asensio-Herrero N, Cristóbal-Azkarate J, Dias PAD, Veà JJ, Rodríguez-Luna E (2007) Foraging habits of *Alouatta palliata mexicana* in three forest fragments. Folia Primatol 78:141–153
- Barrueta T (2003) Población y dieta del mono aullador negro (*Alouatta pigra*) en El Tormento, Campeche, México. Tesis de Maestría, Colegio de la Frontera Sur, Chetumal, Quintana Roo, México

- Behie AM, Pavelka MSM (2005) The short-term effects of a hurricane on the diet and activity of black howlers (*Alouatta pigra*) in Monkey River, Belize. *Folia Primatol* 76:1–9
- Bicca-Marques JC, Calegaro-Marques C (1994a). A case of geophagy in the black howling monkey (*Alouatta caraya*). *Neotrop Primates* 2:7–9
- Bicca-Marques JC, Calegaro-Marques C (1994b) Feeding behavior of the black howler monkey (*Alouatta caraya*) in a seminatural forest. *Acta Biologica Leopoldensia* 16:69–84
- Bicca-Marques JC, Calegaro-Marques C (1994c) Exotic plant species can serve as staple food sources for wild howler populations. *Folia Primatol* 63:209–211
- Bicca-Marques JC, Muhle CB, Prates HM, de Oliveira SG, Calegaro-Marques C (2009) Habitat impoverishment and egg predation by *Alouatta caraya*. *Int J Primatol* 30:743–748
- Biedzicki de Marques AA (1996) Ecology and behavior of brown howlers in araucaria pine forest, southern Brazil. *Neotrop Primates* 4:90–91
- Blake JG, Guerra J, Mosquera D, Torres R, Loiselle BA, Romo D (2010) Use of mineral licks by white-bellied spider monkeys (*Ateles belzebuth*) and red howler monkeys (*Alouatta seniculus*) in Eastern Ecuador. *Int J Primatol* 31:471–483
- Bonvicino CR (1989) Ecología e comportamento de *Alouatta belzebul* (Primates: Cebidae) na Mata Atlântica. *Revista Nordestina de Biologia* 6:149–179
- Bravo SP, Sallenave A (2003) Foraging behavior and activity patterns of *Alouatta caraya* in the northeastern Argentinean flooded forest. *Int J Primatol* 24:825–846
- Bravo SP, Zunino GE (2000) Germination of seeds from three species dispersed by black howler monkeys (*Alouatta caraya*). *Folia Primatol* 71:342–345
- Bravo-Xicoténcatl R. 2003. Patrón diario de actividad comparado en dos tropas de mono aullador (*Alouatta palliata mexicana*) que viven en un mismo fragmento de selva en Los Tuxtlas, Veracruz. BSc Thesis, Universidad Autónoma de Puebla, Puebla

Braza F, Alvarez F, Azcarate T (1983) Feeding habits of the red howler monkeys (*Alouatta seniculus*) in the Llanos of Venezuela. J Mammal 47:205–215

Bridgett GR (2006) The effects of fruit availability and abundance on the diet and ranging behaviour of the black howler monkeys (*Alouatta pigra*) of Monkey River, Belize. MA Thesis, University of Calgary, Calgary, Alberta

Camargo CC, Porfírio S, Rylands AB, Langguth A (2008) Variação sazonal e longitudinal nos padrões de comportamento em uma população de *Alouatta belzebul* (Primates: Atelidae) do Nordeste brasileiro. In: Ferrari SF, Rimoli J (eds) A Primatologia no Brasil 9. Sociedade Brasileira de Primatologia, Biología Geral e Experimental, Aracaju – UFS, pp. 192–201

Carpenter CR (1934) A field study on the behaviour and social relations of howler monkeys. Comp Psychol Monogr 10:1–168

Chapman C (1987a) Flexibility in diets of three species of Costa Rican primates. Folia Primatol 49:80–105

Chapman CA (1987b) Foraging strategies, patch use, and constraints on group size in three species of Costa Rican primates. PhD Thesis, The University of Alberta, Edmonton

Chapman CA (1988) Patterns of foraging and range use by three species of neotropical primates. Primates 29:177–194

Chiarello AG (1993) Activity pattern of the brown howler monkey *Alouatta fusca*, Geoffroy 1812, in a forest fragment of southeastern Brazil. Primates 34:289–293

Chiarello AG (1994) Diet of the brown howler monkey *Alouatta fusca* in a semi-deciduous forest fragment of southeastern Brazil. Primates 35:24–34

Chitolina OP, Sander M (1981) Contribuição ao conhecimento da alimentação de *Alouatta guariba clamitans* Cabrera, 1940 em habitat natural no Rio Grande do Sul (Cebidae, Alouattinae). Iheringia 59:37–44

Coelho AM, Bramblett CA, Quick LB, Bramblett SS (1976) Resource availability and population density in primates: A socio-bioenergetic analysis of the energy budgets of Guatemalan howler and spider monkeys. Primates 17:63–80

Coyohua-Fuentes A (2011) La dieta de los monos aulladores negros (*Alouatta pigra*) en el estado de Campeche: descripción general y variaciones en función del grado de conservación del hábitat. BSc thesis, Universidad Veracruzana, Xalapa

Cristóbal-Azkarate J, Arroyo-Rodríguez V (2007) Diet and activity pattern of howler monkeys (*Alouatta palliata*) in Los Tuxtlas, Mexico: Effects of habitat fragmentation and implications for conservation. Am J Primatol 69:1013–1029

Cuende-Fanton C (2010) Comparación de presupuestos de tiempo, uso del espacio y dieta entre dos grupos de monos aulladores (*Alouatta palliata*) en diferentes condiciones ambientales en Los Tuxtlas, Veracruz, México. MSc thesis, Instituto de Neuroetología, Universidad Veracruzana, Xalapa

Cunha AS (1994) Aspectos sócio-ecológicos de um grupo de bugios (*Alouatta fusca clamitans*) do Parque Estadual Itapuã, RS. MSc thesis, Universidade Federal do Rio Grande do Sul, Porto Alegre

Damé DV (2006) Aspectos ecológicos de um grupo de *Alouatta clamitans* Cabrera, 1940 (Primates, Atelidae) em remanescente de mata Atlântica no município de Camaquã, Rio Grande do Sul. BSc thesis, Universidade Federal de Pelotas, Pelotas, Rio Grande do Sul

de Souza LL, Ferrari SF, Costa ML, Kern DC (2002) Geophagy as a correlate of folivory in red-handed howler monkeys (*Alouatta belzebul*) from eastern Brazilian Amazonia. J Chem Ecol 28:1613–1621

de Thoisy B, Richard-Hansen C (1997) Diet and social behaviour changes in a red howler monkey (*Alouatta seniculus*) troop in a highly degraded rain forest. Folia Primatol 68:357–361

Devos C (1999) Écologie, structure sociale, et différences intersexuelles d'une population de singes hurleurs mantelés (*Alouatta palliata*) sur l'ile d'Ometepe, (Nicaragua): synthèse des résultats d'une recherche conduite en 1999. Cahiers D'Ethologie 20:63–84

Dunn JC, Cristóbal-Azkarate J, Veà JJ (2009) Differences in diet and activity pattern between two groups of *Alouatta palliata* associated with the availability of big trees and fruit of top food taxa. Am J Primatol 71:654–662

Dunn JC, Cristóbal-Azkarate J, Veà JJ (2010) Seasonal variations in the diet and feeding effort of two groups of howlers in different sized forest fragments. *Int J Primatol* 31:887–903

Estrada A (1984) Resource use by howler monkeys (*Alouatta palliata*) in the rain forest of Los Tuxtlas, Veracruz, Mexico. *Int J Primatol* 5:105–131

Estrada A, Coates-Estrada R (1984) Fruit eating and seed dispersal by howling monkeys (*Alouatta palliata*) in the tropical rain forest of Los Tuxtlas, Veracruz, Mexico. *Am J Primatol* 6:77–91

Estrada A, Coates-Estrada R, Yanes CV, Orozco-Segovia A (1984) Comparison of frugivory by howling monkeys (*Alouatta palliata*) and Bats (*Artibeus jamaicensis*) in the tropical rain forest of Los Tuxtlas, Mexico. *Am J Primatol* 7:3–13

Fialho MDS (2000) Ecología do *Alouatta fusca* em floresta de encosta e de restinga no sul do Brasil. MSc dissertation, Instituto de Biologia, Universidade Estadual de Campinas, Campinas, São Paulo

Fortes VB (2008) Ecologia e comportamento do bugio ruivo (*Alouatta guariba clamitans* Cabrera, 1940) em fragmentos florestais na depressão central do Rio Grande do Sul, Brasil. PhD thesis, Pontifícia Universidade Católica do Rio Grande do Sul, Porto Alegre, RS

Fuentes E, Estrada A, Franco B, Magaña M, Decena Y, Muñoz D, García del Valle Y (2003) Reporte preliminar sobre el uso de recursos alimenticios por una tropa de monos aulladores, *Alouatta palliata*, en el parque La Venta, Tabasco, México. *Neotrop Primates* 11:24–29

Galetti M, Pedroni F, Morellato LPC (1994) Diet of the brown howler monkey *Alouatta fusca* in a forest fragment in south-eastern Brazil. *Mammalia* 58:111–118

García del Valle Y, Muñoz D, Magaña M, Estrada A, Franco B (2001) Uso de plantas como alimento por monos aulladores, *Alouatta palliata*, en el Parque Yumká, Tabasco, México. *Neotrop Primates* 9:112–118

García-Orduña F (2002) Comparación de las estrategias de forrajeo de *Ateles geoffroyi vellerosus* y *Alouatta palliata mexicana*, en un fragmento de selva de la Sierra de Santa

Marta, Veracruz. MSc thesis, Instituto de Neuroetología Universidad Veracruzana, Xalapa

Gaspar DA (1997) Ecology and behavior of the brown howling monkey, *Alouatta fusca*. Neotrop Primates 5:80–81

Gaulin SJC, Gaulin CK (1982) Behavioral ecology of *Alouatta seniculus* in Andean cloud forest. Int J Primatol 3:1–31

Giraldo P, Gómez-Posada C, Martín J, Kattan GH (2007) Resource use and seed dispersal by red howler monkeys (*Alouatta seniculus*) in a Colombian Andean forest. Neotrop Primates 14:55–64

Giudice AM, Mudry M (2000) Drinking behavior in the black howler monkey (*Alouatta caraya*). Zoocriaderos 3:11–19

Glander KE (1978) Drinking from arboreal water sources by mantled howling monkeys (*Alouatta palliata* Gray). Folia Primatol 29:206–217

Glander KE (1981) Feeding patterns in mantled howling monkeys. In: Kamil AC, Sargent TD (eds) Foraging behavior: ecological, ethological and psychological approaches. Garland Press, New York, pp. 231–257

González-Picazo H, Estrada A, Ortiz-Martínez T (2001) Consistencias y variaciones en el uso de recursos alimentarios utilizados por una tropa de monos aulladores (*Alouatta palliata*) y deterioro del hábitat en Los Tuxtlas, Veracruz, México. Universidad y Ciencia 17:27–36

Guillotin M, Dubost G, Sabatier D (1994) Food choice and food competition among the three major primate species of French Guiana. J Zool 233:551–579

Guzzo GB (2009) Ecologia e comportamento de *Alouatta guariba clamitans* Cabrera, 1940, em um fragmento de mata de araucária na Serra Gaúcha. MSc thesis, Pontifícia Universidade Católica do Rio Grande do Sul, Porto Alegre, RS

Hladik CM, Hladik A (1969) Rapports trophiques entre végétation et primates dans la forêt de Barro Colorado (Panama). La Terre et al Vie 23:25–117

Izawa K (1975) Foods and feeding behaviour of monkeys in the upper Amazon basin. *Primates* 16:295–316

Jiménez-Huerta J (1992) Distribución y abundancia del recurso alimenticio en un paisaje fragmentado de selva alta perennifolia y su uso por *Ateles* y *Alouatta* en el ejido de Magallanes (Municipio de Soteapan, Veracruz). BSc thesis, Universidad Veracruzana, Xalapa

Juan S, Estrada A, Coates-Estrada R (2000) Contrastes y similitudes en el uso de recursos y patrón general de actividades en tropas de monos aulladores (*Alouatta palliata*) en fragmentos de selva en Los Tuxtlas, México. *Neotrop Primates* 8:131–135

Juan S, Ortiz-Martínez T, Estrada A, Coates-Estrada R (1999) Uso de plantas como alimento por *Alouatta palliata* en un fragmento de selva en Los Tuxtlas, México. *Neotrop Primates* 7:8–11

Julliot C (1996) Fruit choice by red howler monkeys (*Alouatta seniculus*) in a tropical rain forest. *Am J Primatol* 40:261–282

Julliot C, Sabatier D (1993) Diet of the red howler monkey (*Alouatta seniculus*) in French Guiana. *Int J Primatol* 14:527–550

Koch F, Bicca-Marques JC (2007) Padrão de atividades e dieta de *Alouatta guariba clamitans* Cabrera, 1940: uma análise sexo-etária. In: Bicca-Marques JC (ed) A primatologia no Brasil 10. Sociedade Brasileira de Primatologia, Porto Alegre, RS, pp. 353–361

Kuhlmann M (1975) Adenda alimentar dos bugios. *Silvicultura* 9:57–62

Larose F (1996) Foraging strategies, group size, and food competition in the mantled howler monkey, *Alouatta palliata*. PhD thesis, University of Alberta, Edmonton

Limeira VL (1997) Behavioral ecology of *Alouatta fusca clamitans* in a degraded Atlantic forest fragment in Rio de Janeiro. *Neotrop Primates* 5:116–117

Lindberg SM, Santini MEL (1984) A reintrodução do bugio preto (*Alouatta caraya*, Humboldt, 1812 - Cebidae), no Parque Nacional Brasília. *Brasil Florestal* 57:35–53

- Lopez GO, Terborgh J, Ceballos N (2005) Food selection by a hyperdense population of red howler monkeys (*Alouatta seniculus*). *J Trop Ecol* 21:445–450
- Loudon J (2000) Diet, activity and ranging behavior of *Alouatta pigra* in Monkey River, Belize. MA thesis, University of Calgary, Calgary, Alberta
- Ludwig G, Aguiar LM, Svoboda WK, Hilst CLS, Navarro IT, Vitule JRS, Passos FC (2008) Comparison of the diet of *Alouatta caraya* (Primates: Atelidae) between a riparian island and mainland on upper Parana River, southern Brazil. *Rev Bras Zool* 25:419–426
- Lunardelli MC (2000) Padrões de atividade e efeitos de compostos fenólicos na ecologia alimentar de um grupo de bugios-ruivos (*Alouatta fusca*) no sudeste brasileiro. MSc thesis, Universidade de São Paulo, São Paulo
- Marsh LK (1999) Ecological effect of the black howler monkey (*Alouatta pigra*) on fragmented forests in the Community Baboon Sanctuary, Belize. PhD thesis, Washington University, St. Louis
- Marsh LK, Loiselle BA (2003) Recruitment of black howler fruit trees in fragmented forests of northern Belize. *Int J Primatol* 24:65–86
- Martínez-Esquivel LM (2010) Comparación de hábitos alimentarios y su relación con las infecciones parasíticas en los monos congo (*Alouatta palliata*), de Chomes y Palo Verde, Costa Rica. BSc thesis, Universidad de Costa Rica, San José
- Martins CS (1997) Uso de habitat pelo bugio, *Alouatta fusca clamitans*, em um fragmento florestal em Lençóis Paulista. MSc thesis, Universidade Estadual de Campinas, Instituto de Biologia, São Paulo
- Martins MM (2008) Fruit diet of *Alouatta guariba* and *Brachyteles arachnoides* in Southeastern Brazil: comparison of fruit type, color, and seed size. *Primates* 49:1–8
- Martins MM (2009) Lianas as a food resource for brown howlers (*Alouatta guariba*) and southern muriquis (*Brachyteles arachnoides*) in a forest fragment. *Anim Biodiver Conserv* 32:51–58

Mendes SL (1989) Estudo ecológico de *Alouatta fusca* (Primates: Cebidae) na Estação de Caratinga, MG. Revista Nordestina de Biologia 6:71–104

Milton K (1980) The foraging strategy of howler monkeys; a study of primate economics. Columbia University Press, New York

Miranda JMD, Moro-Rios RF, Bernardi IP, Passos FC (2005) Formas não usuais para a obtenção de água por *Alouatta guariba clamitans* em ambiente de floresta com Araucária no sul do Brasil. Neotrop Primates 13:21–23

Miranda JMD, Passos FC (2004) Hábito alimentar de *Alouatta guariba* (Humbolt) (Primates, Atelidae) em floresta de Araucária, Paraná, Brasil. Rev Bras Zool 21:821–826

Mittermeier RA, van Roosmalen MGM (1981) Preliminary observations on habitat utilization and diet in eight Surinam monkeys. Folia Primatol 36:1–39

Muhle CB (2008) Estratégias adaptativas de um grupo de bugios-pretos, *Alouatta caraya* (Humboldt, 1812), em um bosque dominado por eucaliptos. MSc thesis, Pontifícia Universidade Católica do Rio Grande do Sul, Porto Alegre, RS

Muñoz D, García del Valle Y, Franco B, Estrada A, Magaña M (2002) Estudio del patrón de actividad general de monos aulladores (*Alouatta palliata*) en el parque Yumká, Tabasco, México. Neotrop Primates 10:11–17

Neves AM, Rylands AB (1991) Diet of a group of howling monkeys, *Alouatta seniculus*, in an isolated forest patch in central Amazonia. In: Rylands AB, Bernardes AT (eds) A primatología no Brasil. Universidade Federal de Juiz de Fora, Minas Gerais, pp. 263–274

Ortiz-Martínez T, Juan-Solano S, Estrada A, Coates-Estrada R (1999) Patrones de actividad de *Alouatta palliata* en un fragmento de selva en Los Tuxtlas, Mexico. Neotrop Primates 7:80–88

Palma AC, Veléz A, Gómez-Posada C, López H, Zárate DA, Stevenson PR (2011) Use of space, activity patterns, and foraging behavior of red howler monkeys (*Alouatta seniculus*) in an Andean forest fragment in Colombia. Am J Primatol 73:1062–1071

- Pavé RE, Peker S, Raño M, Orjuela CR, Zunino GE, Kowalewski MM (2009) Nectar feeding on an exotic tree (*Grevillea robusta*) by *Alouatta caraya* and its possible role in flower pollination. *Neotrop Primates* 16:61–64
- Pavelka MSM, Knopff K (2004) Diet and activity in black howler monkeys (*Alouatta pigra*) in southern Belize: does degree of frugivory influence activity level? *Primates* 45:105–111
- Pereira TS (2008) Ecologia cognitiva e forrageamento de *Alouatta guariba clamitans* Cabrera, 1940: os bugios ruivos possuem mapas mentais? MSc thesis, Pontifícia Universidade Católica do Rio Grande do Sul, Porto Alegre, RS
- Perez DM (1997) Estudo ecológico do bugiô ruivo em uma floresta com araucária do sul do Brasil (*Alouatta fusca*, Ihering 1914 - Primates, Atelidae). MSc thesis, Universidade de São Paulo, São Paulo
- Pinto ACB, Azevedo-Ramos C, Carvalho O (2003) Activity patterns and diet of the howler monkey *Alouatta belzebul* in areas of logged and unlogged forest in Eastern Amazonia. *Anim Biodiver Conserv* 26:39–49
- Pinto LP (2002) Dieta padrão de atividades e área de vida de *Alouatta belzebel discolor* (Primates, Atelidae) em Paranaíta, Norte de Mato Grosso. MSc thesis, Instituto de Biología, Universidade Estadual de Campinas, São Paulo
- Pinto LP, Setz EZ (2004) Diet of *Alouatta belzebul discolor* in an Amazonian rain forest of northern Mato Grosso State, Brazil. *Int J Primatol* 25:1197–1211
- Pozo-Montuy G, Serio-Silva JC (2006) Comportamiento alimentario de monos aulladores negros (*Alouatta pigra* Lawrence, Cebidae) en hábitat fragmentado en Balancán, Tabasco, México. *Acta Zoológica Mexicana* 22:53–66
- Pozo-Montuy G, Serio-Silva JC (2007) Movement and resource use by a group of *Alouatta pigra* in a forest fragment in Balancán, México. *Primates* 48:102–107
- Prates HM (2007) Ecologia e comportamento de um grupo de bugios pretos (*Alouatta caraya*) habitante de um pomar em Alegrete, RS, Brasil. MSc thesis, Pontifícia Universidade Católica do Rio Grande do Sul, Porto Alegre, RS

- Prates HM, Bicca-Marques JC (2008) Age-sex analysis of activity budget, diet, and positional behavior in *Alouatta caraya* in an orchard forest. *Int J Primatol* 29:703–715
- Queiroz HL (1995) Preguiças e guaribas: os mamíferos folívoros arborícolas do Mamirauá. Sociedade Civil Mamirauá & CNPq, Brasilia
- Rímoli AO, Valdivino EM, Rímoli J, Ferrari SF (2008) Behavior patterns of a group of black howler monkeys *Alouatta caraya* (Humboldt, 1812) in a forest fragment in Terenos, Mato Grosso do Sul: a seasonal analysis. In: Rímoli J, Ferrari SF (eds) A primatologia no Brasil 9. Sociedade Brasileira de Primatologia, Biología Geral e Experimental – UFS, Aracaju
- Rivera A, Calmé S (2006) Forest fragmentation and its effects on the feeding ecology of black howlers (*Alouatta pigra*) from the Calakmul area in Mexico. In: Estrada A, Garber PA, Pavelka MSM, Luecke L (eds) New perspectives in the study of Mesoamerican primates: Distribution, ecology, behavior, and conservation. Springer, New York, pp. 189–213
- Rizzo KA (2004) The sociecology of *Alouatta pigra* at the Calakmul archeological zone and biosphere reserve, Mexico. PhD thesis, University of Illinois, Chicago
- Rodríguez-Luna E, Domínguez-Domínguez LE, Morales-Mávil J, Martínez-Morales MM (2003) Foraging strategy changes in an *Alouatta palliata mexicana* troop released on an island. In: Marsh LK (ed) Primates in fragments: ecology and conservation. Kluwer Academic/Plenum Publishers, New York, pp. 229–250
- Rumiz DI, Zunino GE, Obregonzo ML, Ruiz JC (1986) *Alouatta caraya*: habitat and resource utilization in northern Argentina. In: Taub DM, King AW (eds) Current perspectives in primate social dynamics. Van Nostrand Reinhold Company, New York, pp. 175–193
- Schlichte HJ (1978) A preliminary report on the habitat utilization of a group of howler monkeys (*Alouatta villosa pigra*) in the National Park of Tikal, Guatemala. In: Montgomery GG (ed) Ecology of arboreal folivores. Smithsonian Institution Press, Washington, pp. 551–559

Sedden-González A, Rodríguez-Luna E (2010) Responses of a translocated howler monkey *Alouatta palliata* group to new environmental conditions. *Endangered Species Research* 12:25–30

Serio-Silva JC (1995) Patrón diario de actividades y hábitos alimenticios de *Alouatta palliata* en semilibertad. In: Rodríguez-Luna E, Cortés-Ortiz L, Contreras JM (eds) *Estudios primatológicos en México, Vol. II*. Universidad Veracruzana, Xalapa, pp. 149–171

Serio-Silva JC, Rico-Gray V, Hernández-Salazar LT, Espinosa-Gomez R (2002) The role of *Ficus* (Moraceae) in the diet and nutrition of a troop of Mexican howler monkeys, *Alouatta palliata mexicana*, released on an island in southern Veracruz, Mexico. *J Trop Ecol* 18:913–928

Silva EC (1981) A preliminary survey of brown howler monkeys (*Alouatta fusca*) at the Cantareira Reserve (São Paulo, Brazil). *Rev Bras Biol* 41:897–909

Silver SC, Marsh LK (2003) Dietary flexibility, behavioral plasticity, and survival in fragments: lessons from translocated howlers. In: Marsh LK (ed) *Primates in fragments: ecology and conservation*. Kluwer Academic/Plenum Publishers, New York, pp. 251–265

Silver SC, Ostro LET, Yeager CP, Horwich RH (1998) Feeding ecology of the black howler monkey (*Alouatta pigra*) in northern Belize. *Am J Primatol* 45:263–279

Simmen B, Sabatier D (1996) Diets of some French Guianan primates: composition and food choices. *Int J Primatol* 17:661–693

Smith CC (1977) Feeding behaviour and social organization in howler monkeys. In: Clutton-Brock TH (ed) *Primate ecology*. Academic Press, London, pp. 97–126

Snarr KA (2006) Life in a lowland wet forest fragment on the north coast of Honduras: the mantled howlers (*Alouatta palliata*) of Cuero y Salado Wildlife Refuge. PhD thesis, University of Toronto, Toronto

Soini P (1992) Ecología del coto mono (*Alouatta seniculus*, Cebidae) en el Río Pacaya, Reserva Nacional Pacaya-Samiria, Peru. *Folia Amazonica* 4:117–134

- Steinmetz S (2000) Ecologia e o comportamento do bugio (*Alouatta fusca clamitans*, Atelidae - Primates) no Parque Estadual Intervales. MSc thesis, Universidade de São Paulo, São Paulo
- Steinmetz S (2001) Drinking by howler monkeys (*Alouatta fusca*) and its seasonality at the Intervales State Park, São Paulo, Brazil. *Neotrop Primates* 9:111–112
- Stevenson PR, Castellanos MC, Pizarro JC, Garavito M (2002) Effects of seed dispersal by three ateline monkey species on seed germination at Tinigua National Park, Colombia. *Int J Primatol* 23:1187–1204
- Stevenson PR, Quiñones MJ, Ahumada JA (2000) Influence of fruit availability on ecological overlap among four Neotropical primates at Tinigua National Park, Colombia. *Biotropica* 32:533–544
- Stoner KE (1996) Habitat selection and seasonal patterns of activity and foraging of mantled howling monkeys (*Alouatta palliata*) in northeastern Costa Rica. *Int J Primatol* 17:1–30
- Tomblin DC, Cranford JA (1994) Ecological niche differences between *Alouatta palliata* and *Cebus capucinus* comparing feeding modes, branch size use, and diet. *Primates* 35:265–274
- Welker BJ (2004) Proximate mechanisms governing feeding behavior and selectivity in mantled howler monkeys, *Alouatta palliata*. Ph.D. thesis, State University of New York, Buffalo
- Williams-Guillén K (2003) The behavioral ecology of mantled howling monkeys (*Alouatta palliata*) living in a Nicaraguan shade coffee plantation. New York University, New York
- Young AL (1983) Preliminary observations on the ecology and behavior of the muriqui and brown howler monkey. Bachelor's thesis, Harvard University, Cambridge, Massachusetts
- Zunino GE (1986) Algunos aspectos de la ecología y etología del mono aullador negro (*Alouatta caraya*) en hábitat fragmentados. Ph.D thesis, Universidad de Buenos Aires, Buenos Aires

Zunino GE (1989) Hábitat, dieta y actividad del mono aullador negro (*Alouatta caraya*) en el noroeste de la Argentina. Boletín Primatológico Latinoamericano 1:47–97