Analysis of translation-invariant operators via the Fourier transform of the reproducing kernel

Egor Maximenko, joint results with

Gerardo Ramos-Vazquez, Crispin Herrera-Yañez, and Alejandro Hernández Arteaga

Instituto Politécnico Nacional Escuela Superior de Física y Matemáticas, México

2022-09-27

Operator Theory and Function Spaces, Xalapa, Mexico

Outline



2 Separately radial operators/Bergman space



Vertical operators/poly-Fock space 000000000000

(1) Scheme for domains $G \times Y$

2 Separately radial operators/Bergman space

③ Vertical operators/poly-Fock space

A general problem (we cannot solve it)

Let

- X be a set,
- H be a reproducing kernel Hilbert space over X,
- G be a locally compact group,

 $\tau: G \to \operatorname{Sym}(X)$ be a group action, $(\rho(g))_{g \in G}, \quad \rho(g) f := f \circ \tau(g^{-1})$ be a unitary representation of G in H.

Problem: describe the W*-algebra defined as the centralizer of ρ ,

$$\mathcal{C}(
ho) \ \coloneqq \ \Big\{ S \in \mathcal{B}(\mathcal{H}) \colon \quad orall g \in \mathcal{G} \quad S \,
ho(g) =
ho(g) \, S \Big\}.$$

A general idea

Apply the Fourier transform to the reproducing kernel along the orbits of the group action:

$$\int_G \mathsf{K}_{\mathsf{z}}(\tau(\mathsf{g})(\mathsf{w}))\,\psi(\mathsf{g})^*\,\mathrm{d}\nu_G(\mathsf{g}),\qquad\psi\in\mathsf{irreducible representations of }\,G.$$

We hope that the obtained operator-valued function is useful to describe $C(\rho)$.

Our scheme for type-type domains $G \times Y$

 Crispin Herrera-Yañez, Egor A. Maximenko, Gerardo Ramos-Vazquez (2022): Translation-invariant operators in reproducing kernel Hilbert spaces. Integral Equ. Oper. Theory. DOI: 10.1007/s00020-022-02705-4.

Our paper is inspired by various works of Vasilevski and other mathematicians.

Nikolai L. Vasilevski (1999):

On Bergman-Toeplitz operators with commutative symbol algebras. Integral Equ. Oper. Theory. DOI: 10.1007/BF01332495.

Our assumptions

- $X = G \times Y$,
- G is an abelian locally compact group, metrizable, and σ -compact,
- ν is a Haar measure on G,
- (Y, λ) is a σ -finite measure space,
- $L^2(G \times Y)$ is separable,
- $H \leq L^2(G \times Y)$,
- (*H* is an RKHS); we denote the RK by $(K_{x,y})_{x \in G, y \in Y}$,

Our assumptions

• G acts in $G \times Y$ by

$$au_{G \times Y}(g)$$
: $(x, y) \mapsto (g + x, y),$

• $\rho_{G \times Y}$ acts in $L^2(G \times Y)$

$$(\rho_{G\times Y}(a)f)(x,y) \coloneqq f(x-a,y),$$

• *H* is invariant under $\rho_{G \times Y}$,

•
$$\forall y \in Y \quad \sup_{v \in Y} \int_{G} |K_{(0,y)}(u,v)| d\nu(u) < +\infty.$$

Criterion that H is shift-invariant

P := the orthogonal projection in $L^2(G \times Y)$ whose image is H.

Proposition

The following conditions are equivalent.

(a)
$$\rho_{G \times Y}(a)(H) \subseteq H$$
 for every a in G .

(b)
$$P\rho_{G\times Y}(a) = \rho_{G\times Y}(a)P$$
 for every *a* in *G*.

(c) $K_{x,y}(u,v) = K_{0,y}(u-x,v)$ for every x, y in G and every y, v in Y.

(d) $\rho_{G \times Y}(a)K_{x,y} = K_{a+x,y}$ for every a, x in G and every y in Y.

Let $\rho_H(a)$: $H \to H$ be the compression of $\rho_{G \times Y}(a)$.

Vertical operators/poly-Fock space 000000000000

Decomposition of H

$$\widehat{P} \coloneqq (F \otimes I)P(F \otimes I)^*, \qquad \widehat{H} \coloneqq (F \otimes I)(H).$$
 $\widehat{P} = \int_{\widehat{G}}^{\oplus} \widehat{P}_{\xi} d\widehat{\nu}(\xi).$

For each ξ in \widehat{G} ,

$$\widehat{H}_{\xi} \coloneqq \widehat{P}_{\xi}(L^{2}(Y)).$$
 $\Omega \coloneqq \{\xi \in \widehat{G} \colon \dim(\widehat{H}_{\xi}) > 0\}.$
 $\widehat{H} = \int_{\Omega}^{\oplus} \widehat{H}_{\xi} d\widehat{\nu}(\xi).$

Scheme for domains $G \times Y$

Separately radial operators/Bergman space

Vertical operators/poly-Fock space 000000000000

Decomposition of $\mathcal{V} \coloneqq \mathcal{C}(\rho_H)$

Let $\Phi \colon H \to \widehat{H}$ be the compression of $F \otimes I$.

Theorem

$$\Phi \, \mathcal{V} \, \Phi^* = \int_\Omega^\oplus \mathcal{B}(\widehat{H}_\xi) \, \mathrm{d} \widehat{
u}(\xi).$$



Scheme for domains $G \times Y$

Separately radial operators/Bergman space

Vertical operators/poly-Fock space 000000000000

Constructive description of the fibers \widehat{H}_{ξ}

$$L_{\cdot,y} := (F \otimes I) K_{0,y}, \qquad \text{i.e.}, \qquad L_{\xi,y}(v) := \int_G K_{(0,y)}(u,v) \overline{\xi(u)} \, \mathrm{d}\nu(u).$$

Theorem

For every ξ in Ω , the family $(L_{\xi,y})_{y\in Y}$ is the reproducing kernel of \widehat{H}_{ξ} .

Idea of the proof: convolution theorem + Fubini + Moore-Aronszajn theorem.

Constructive criterion for the commutativity of $\ensuremath{\mathcal{V}}$

Theorem

The following conditions are equivalent.

(a) \mathcal{V} is commutative.

(b)
$$d_{\xi} \coloneqq \operatorname{dim}(\widehat{H}_{\xi}) = 1$$
 for every ξ in Ω .

(c) $|L_{\xi,y}(v)|^2 = L_{\xi,y}(y)L_{\xi,v}(v)$ for every ξ in Ω and every y, v in Y.

(d) There exists a family $(q_{\xi})_{\xi \in \Omega}$ in $L^2(Y)$ such that the function $(\xi, v) \mapsto q_{\xi}(v)$ is measurable, $\widehat{H}_{\xi} = \mathbb{C}q_{\xi}$, $||q_{\xi}|| = 1$, and

$$L_{\xi,y}(v) = \overline{q_{\xi}(y)}q_{\xi}(v) \qquad (\xi \in \Omega, \ y, v \in Y).$$

Vertical operators/poly-Fock space

Isometric isomorphism $R: H \to L^2(\Omega)$ in the commutative case

Suppose that dim $(\hat{H}_{\xi}) = 1$, i.e., we have a family $(q_{\xi})_{\xi \in \Omega}$ such that $\hat{H}_{\xi} = \mathbb{C}q_{\xi} \text{ y } ||q_{\xi}|| = 1$.

 $(Rf)(\xi) \coloneqq \langle (\Phi f)(\xi, \cdot), q_{\xi} \rangle_{L^{2}(Y)}.$



Vertical operators/poly-Fock space 000000000000

Diagonalization of translation-invariant operators in the case $d_{\xi}=1$

Proposition

Suppose that dim $(\widehat{H}_{\xi}) = 1$ for every ξ in Ω . Entonces $\mathcal{V} \cong L^{\infty}(\Omega)$.



Vertical operators/poly-Fock space 000000000000

The case of finite-dimensional fibers

Suppose that

$$orall \xi \in \Omega \qquad d_{\xi} := \dim(\widehat{H}_{\xi}) < +\infty.$$

Let $(q_{j,\xi})_{j\in\mathbb{N},\xi\in\Omega}$ be a measurable basis family for the spaces \widehat{H}_{ξ} .

$$L_{\xi,y}(v) = \sum_{j=1}^{d_{\xi}} \overline{q_{j,\xi}(y)} \, q_{j,\xi}(v).$$

Then

$$\Phi H = \widehat{H} = \int_{\Omega}^{\oplus} \widehat{H}_{\xi} \, \mathrm{d}\widehat{\nu}(\xi) \cong \int_{\Omega}^{\oplus} \mathbb{C}^{d_{\xi}} \, \mathrm{d}\widehat{\nu}(\xi).$$

Vertical operators/poly-Fock space

From translation-invariant operators to matrix families

$$R \colon H o \int_{\Omega}^{\oplus} \mathbb{C}^{d_{\xi}} \operatorname{d} \widehat{
u}(\xi), \qquad (Rf)(\xi) \coloneqq \left[\langle (\Phi f)(\xi, \cdot), \ q_{j,\xi}
angle_{L^2(Y)}
ight]_{j=1}^{d_{\xi}}$$



Vertical operators/poly-Fock space

Matrix families corresponding to Toeplitz operators with translation-invariant generating symbols

Corollary

Let $\psi \in L^{\infty}(Y)$,

$$\varphi(\mathbf{x},\mathbf{y})=\psi(\mathbf{y}).$$

Then $T_{arphi} \in \mathcal{V}$, $RT_{arphi}R^* = M_{\gamma_{\psi}}$,

$$\gamma_{\psi}(\xi) \coloneqq \left[\int_{\mathbf{Y}} \psi(\mathbf{v}) \overline{q_{j,\xi}(\mathbf{v})} q_{k,\xi}(\mathbf{v}) \,\mathrm{d}\lambda(\mathbf{v})
ight]_{j,k=1}^{d_{\xi}}$$



Vertical operators/poly-Fock space 000000000000



2 Separately radial operators/Bergman space

③ Vertical operators/poly-Fock space

Vertical operators/poly-Fock space

Separately radial case on the Bergman space

Raul Quiroga-Barranco, Nikolai Vasilevski (2007):

Commutative C*-algebras of Toeplitz operators on the unit ball, I. Bargmann-type transforms and spectral representations of Toeplitz operators. Integral Equ. Oper. Theory. DOI: 10.1007/s00020-007-1537-6.

They worked with maximal abelian subgroups of the Möbius group and diagonalized Toeplitz operators with group-invariant symbols.

Jointly with Alejandro Hernández Arteaga,

we studied three of these cases using the scheme above.

In this talk, we will see the separately radial (= quasi-radial) case.

Vertical operators/poly-Fock space

$\mathcal{A}^2(\mathbb{B}_n, \mu_{n,\alpha})$, the analytic Bergman space

$$\mathbb{B}_n \coloneqq \{z \in \mathbb{C}^n \colon |z| < 1\}.$$

$$\mathrm{d}\mu_{n,\alpha}(z) = c_{n,\alpha} \left(1 - |z|^2\right)^{\alpha} \mathrm{d}\mu_{2n}(z), \qquad c_{n,\alpha} = \frac{\Gamma(n+\alpha+1)}{\pi^n \, \Gamma(\alpha+1)}.$$

 $\mathcal{A}^2 = \mathcal{A}^2(\mathbb{B}_n, \mu_{n,\alpha}) \coloneqq$ holomorphic functions belonging to $L^2(\mathbb{B}_n, \mu_{n,\alpha})$.

Orthonormal basis:
$$b_j(z) = \sqrt{rac{\Gamma(n+|j|+lpha+1)}{j!\,\Gamma(n+lpha+1)}}\,z^j, \qquad j\in\mathbb{N}_0^n.$$

Reproducing kernel of \mathcal{A}^2 : $\mathcal{K}_z^{\mathcal{A}^2}(w) = \frac{1}{(1 - \langle w, z \rangle)^{n+1+\alpha}}.$

Vertical operators/poly-Fock space

Group $\mathbb{R}_{2\pi}^n$ and its dual group

$$G \coloneqq \mathbb{R}_{2\pi}^n \cong \mathbb{T}^n$$
, where $\mathbb{R}_{2\pi} \coloneqq \mathbb{R}/(2\pi\mathbb{Z})$.

 $\nu :=$ the normalized Haar measure on *G*.

$$\int_G f \,\mathrm{d}\nu = \frac{1}{(2\pi)^n} \int_{[0,2\pi)^n} f(g) \,\mathrm{d}\mu_n(g).$$

 $\widehat{G} = \mathbb{Z}^n$ with the counting measure $\widehat{\nu}$.

Pairing between *G* and \widehat{G} :

$$E(u+2\pi\mathbb{Z}^n,\xi)=\mathrm{e}^{\mathrm{i}\,\langle u,\xi\rangle}\,.$$

Vertical operators/poly-Fock space

Rotations acting in \mathcal{A}^2

 $G = \mathbb{R}^n_{2\pi} \cong \mathbb{T}^n.$

Action of *G* on \mathbb{B}_n :

$$\tau_{\rm rot}(g)(z) \coloneqq ({\rm e}^{{\rm i}\,g_1}\,z_1,\ldots,{\rm e}^{{\rm i}\,g_n}\,z_n).$$

Unitary representation of G in \mathbb{B}_n :

$$(\rho_{\mathsf{rot}}(g)f)(z) \coloneqq f(\tau_{\mathsf{rot}}(-g)z) = f(e^{-ig_1}z_1, \dots, e^{-ig_n}z_n).$$

Passing to the polar coordinates

Let Y be the base of \mathbb{B}_n considered as a Reinhard domain:

$$Y=\Big\{y\in [0,+\infty)^n\colon \quad |y|^2<1\Big\}.$$

We consider Y with the Lebesgue measure μ_n .

$$\varphi_{\mathsf{polar}} \colon G \times Y \to \mathbb{B}_n, \qquad \varphi_{\mathsf{polar}}(u, v) = (v_1 e^{i u_1}, \dots, v_n e^{i u_n}).$$

For every f in \mathcal{A}^2 , we define $U_{polar}f \in L^2(G \times Y)$,

$$(U_{\mathsf{polar}}f)(u,v)\coloneqq \sqrt{(2\pi)^n c_{n,lpha} v_1 \cdots v_n} \ (1-|v|^2)^{lpha/2} \ f(arphi_{\mathsf{polar}}(u,v)).$$

 $H \coloneqq U_{\mathsf{polar}}(\mathcal{A}^2). \quad U_{\mathsf{polar}} \colon \mathcal{A}^2 \to H \text{ is an isometric isomophism.}$

Vertical operators/poly-Fock space 000000000000

Transformation of RK by a weighted change of variable



Then the following function is the RK of H_2 :

$$\widetilde{K}_{z}(w) = \overline{p(z)} \, p(w) \, K_{\varphi(z)}(\varphi(w)).$$

Vertical operators/poly-Fock space

Passing to the polar coordinates

The reproducing kernel of H is

$$\mathcal{K}_{x,y}(u,v) = \frac{(2\pi)^n c_{n,\alpha} (1-|y|^2)^{\alpha/2} (1-|v|^2)^{\alpha/2} \prod_{k=1}^n \sqrt{y_k v_k}}{\left(1-\sum_{k=1}^n y_k v_k e^{i(u_k-x_k)}\right)^{n+\alpha+1}}.$$

We see that $K_{x,y}(u, v) = K_{0,y}(u - x, v)$.

Furthermore, U_{polar} intertwines ρ_{rot} with horizontal translations:

$$orall g \in G \qquad U_{ extsf{polar}} \,
ho_{ extsf{rot}}(g) =
ho_{G imes Y}(g) \, U_{ extsf{polar}}.$$

Vertical operators/poly-Fock space 000000000000

U_{polar} intertwines the rotations acting in \mathcal{A}^2 with the horizontal translations acting in H



Vertical operators/poly-Fock space 000000000000

Computation of L

1

 $K_{0,y}(\cdot, v)$ decomposes into the Fourier series:

$$\begin{split} \mathsf{K}_{0,y}(u,v) &= (2\pi)^n \, c_{n,\alpha} \, (1-|v|^2)^{\alpha/2} \, (1-|y|^2)^{\alpha/2} \prod_{k=1}^n \sqrt{v_k y_k} \times \\ & \times \sum_{\xi \in \mathbb{N}_0^n} \frac{\mathsf{\Gamma}(n+|\xi|+\alpha+1)}{\xi! \, \mathsf{\Gamma}(n+\alpha+1)} \, y^{\xi} \, v^{\xi} \, \operatorname{e}^{\operatorname{i} \langle u, \xi \rangle} \, . \end{split}$$

For ξ in \mathbb{N}_0^n , the ξ th Fourier coefficient is $L_{\xi,y}(v) = \overline{q_{\xi}(y)} q_{\xi}(v)$, where

$$q_{\xi}(v) = \sqrt{\frac{2^n \, \Gamma(n+|\xi|+\alpha+1)}{\xi! \, \Gamma(\alpha+1)}} \, (1-|v|^2)^{\alpha/2} \, \prod_{k=1}^n v_k^{\xi_k+\frac{1}{2}}.$$

Vertical operators/poly-Fock space 000000000000

Conclusions

In this example:

$$\Omega = \mathbb{N}_0^n$$
,

$$egin{aligned} &d_{\xi}=1 \mbox{ for } \xi \mbox{ in } \mathbb{N}^n_0, \ &\mathcal{C}(
ho_{
m rot})\cong\mathcal{C}(
ho_H)=\mathcal{V}\cong L^\infty(\mathbb{N}^n_0). \end{aligned}$$

 $\ensuremath{\mathcal{V}}$ is commutative.



Vertical operators/poly-Fock space

The eigenvalues of separately radial Toeplitz operators in $\mathcal{A}^2(\mathbb{B}_n)$

We suppose that $\varphi \in L^{\infty}(Y)$.

$$\begin{split} \gamma_{\varphi}(\xi) &= \int_{Y} \varphi(v) \, |q_{\xi}(v)|^2 \, \mathrm{d}\mu_n(v) \\ &= \frac{\Gamma(n+|\xi|+\alpha+1)}{\xi! \, \Gamma(\alpha+1)} \int_{|t|_1 < 1} \varphi(\sqrt{t}) \, (1-|t|_1)^{\alpha} \, t^{\xi} \, \mathrm{d}\mu_n(t). \end{split}$$

Here $|t|_1 = t_1 + \cdots + t_n$.

This formula coincides with the formula found by Quiroga-Barranco and Vasilevski.

Separately radial operators on the pluriharmonic Bergman space

$$rac{1}{(1-\langle w,z
angle)^{n+1+lpha}}+rac{1}{(1-\langle z,w
angle)^{n+1+lpha}}-1.$$

The analysis of separately radial operators is similar to the analytic case, but

$$\Omega = \mathbb{N}_0^n \cup (-\mathbb{N}_0)^n = \{0, 1, 2, \ldots\}^n \cup \{0, -1, -2, \ldots\}^n.$$

 $\mathcal{V} \cong L^{\infty}(\Omega)$. C*-alg(sep. radial Toeplitz operators) is **not** weakly dense in \mathcal{V} .

- Jingyu Yang, Liu Liu, Yufeng Lu (2013).
- Maribel Loaiza, Carmen Lozano (2014).

Vertical operators/poly-Fock space

Separately radial operators on the pluriharmonic Bergman space

$$\Omega = \mathbb{N}_0^n \cup (-\mathbb{N}_0)^n.$$



Outline



2 Separately radial operators/Bergman space



This is a beginning of a joint work with

Gerardo Ramos Vazquez, Armando Sánchez Nungaray, and Erick Lee Guzmán.

At the moment, we have applied the $G \times Y$ scheme to the vertical operators on the *m*-analytic Fock space $\mathcal{F}_m(\mathbb{C}^1)$. Our results are similar to the following two papers.

Nikolai L. Vasilevski (2000):

Poly-Fock spaces.

Armando Sánchez-Nungaray, Carlos González-Flores,
 Raquiel Rufino López-Martínez, Jorge Luis Arroyo-Neri (2018):
 Toeplitz operators with horizontal symbols acting on the poly-Fock spaces.

Vertical operators/poly-Fock space

Polyanalytic Bargmann–Segal–Fock space

 $\mathcal{F}_m \coloneqq m$ -analytic functions $\mathbb{C} \to \mathbb{C}$, square integrable with weight $e^{-|z|^2}$.

$$\|f\|_{\mathcal{F}_m}\coloneqq \left(rac{1}{\pi}\int_{\mathbb{C}}|f(z)|^2\,\mathrm{e}^{-|z|^2}\,\mathrm{d}\mu_2(z)
ight)^{1/2},$$

Nour eddine Askour, Ahmed Intissar, Zouhaïr Mouayn (1997) computed the reproducing kernel of this space:

$$K_z^{\mathcal{F}_m}(w) = e^{\overline{z}w} L_{m-1}^{(1)}(|w-z|^2).$$

Vertical operators/poly-Fock space

Polyanalytic Steinwart–Hush–Scovel space

 $\mathcal{S}_m \coloneqq m$ -analytic functions $\mathbb{C} \to \mathbb{C}$ such that

$$\|f\|_{\mathcal{S}_m}\coloneqq \left(rac{1}{\pi}\int_{\mathbb{C}}|f(z)|^2\exp\left(-2\operatorname{Im}(z)^2
ight)\mathrm{d}\mu_2(z)
ight)^{1/2}<+\infty.$$

Isometric isomorphism $U_{\mathcal{F}_m}^{\mathcal{S}_m} \colon \mathcal{F}_m o \mathcal{S}_m$,

$$(U_{\mathcal{F}_m}^{\mathcal{S}_m}f)(z)\coloneqq \mathrm{e}^{-z^2/2}\,f(z).$$

Reproducing kernel of S_m :

$$K_z^{\mathcal{S}_m}(w) = e^{-\frac{1}{2}(w-\overline{z})^2} L_{m-1}^{(1)}(|w-z|^2).$$

The original Steinwart–Hush–Scovel space on \mathbb{C}^n

6

Ingo Steinwart, Don Hush, Clint Scovel (2006).

Analytic functions on \mathbb{C}^n such that

$$\frac{2^n\alpha^{2n}}{\pi^n}\int_{\mathbb{C}^n}|f(z)|^2\exp\left(-4\alpha^2\sum_{j=1}^n\operatorname{Im}(z_j)^2\right)\mathrm{d}\mu_{2n}(z)<+\infty.$$

Reproducing kernel:

$$\exp\left(-lpha^2\sum_{j=1}^n(w_j-\overline{z_j})^2
ight).$$

Its restriction to \mathbb{R}^n (the Gaussian kernel) is widely used in machine learning.

Vertical operators/poly-Fock space

"Flattened poly-Fock space"

Isometric isomorphism $U_{\mathcal{F}_m}^{\mathcal{H}_m} \colon \mathcal{F}_m \to \mathcal{H}_m < L^2(\mathbb{R}^2)$,

$$(U_{\mathcal{F}_m}^{\mathcal{H}_m}f)(x,y) \coloneqq \left(\frac{2}{\pi}\right)^{1/4} e^{-\frac{x^2+y^2}{2}-i\,xy}\,f(x+i\,y).$$

Reproducing kernel of \mathcal{H}_m :

$$K_{x,y}(u,v) = \sqrt{\frac{2}{\pi}} e^{-\frac{(u-x)^2 + (v-y)^2}{2} - i(u-x)(v+y)} L_{m-1}^{(1)}((u-x)^2 + (v-y)^2).$$

We see that $K_{x,y}(u,v) = K_{0,y}(u-x,v).$

Vertical operators/poly-Fock space

\mathcal{S}_m and \mathcal{H}_m are "flattened" versions of the poly-Fock space \mathcal{F}_m



Vertical operators/poly-Fock space

Weyl operators and horizontal translations

Unitary representation of \mathbb{R} in \mathcal{F}_m :

$$(
ho_{\mathcal{F}_m}(a)f)(z) \coloneqq f(z-a) e^{az-rac{a^2}{2}}.$$

Unitary representation of \mathbb{R} in \mathcal{S}_m :

$$(\rho_{\mathcal{S}_m}(a)f)(z) := f(z-a).$$

Unitary representation of \mathbb{R} in \mathcal{H}_m :

$$(\rho_{\mathcal{H}_m}(a)f)(x,y) := f(x-a,y).$$

Vertical operators/poly-Fock space

Three Hilbert spaces and corresponding unitary representations of $\mathbb R$



Vertical operators/poly-Fock space

Fourier connection between Laguerre and Hermite functions

$$\frac{1}{\sqrt{2\pi}} \int_{\mathbb{R}} e^{-i u\xi} e^{-\frac{u^2}{2}} L_n(u^2 + a^2) du = \frac{1}{2^n n!} e^{-\frac{\xi^2}{2}} H_n\left(\frac{\xi + a}{\sqrt{2}}\right) H_n\left(\frac{\xi - a}{\sqrt{2}}\right).$$

Equivalently,

$$\frac{1}{\sqrt{2\pi}}\int_{\mathbb{R}} e^{-i u\xi} e^{-\frac{u^2+a^2}{2}} L_n(u^2+a^2) du = \sqrt{\pi} \psi_n\left(\frac{\xi+a}{\sqrt{2}}\right) \psi_n\left(\frac{\xi-a}{\sqrt{2}}\right).$$

Here ψ_n is the *n*th Hermite function: $\psi_n(t) = \frac{1}{\sqrt{2^n n! \sqrt{\pi}}} e^{-t^2/2} H_n(t).$

Serald B. Folland (1989). Harmonic Analysis on Phase Space.

Vertical operators/poly-Fock space

Fourier transform of the reproducing kernel of \mathcal{H}_m

Proposition

For every ξ, y, v in \mathbb{R} ,

$$L_{\xi,y}(v) = \frac{1}{\sqrt{2\pi}} \int_{\mathbb{R}} \mathcal{K}_{0,y}(u,v) e^{-i u\xi} du = \sum_{j=0}^{m-1} \overline{q_{j,\xi}(y)} q_{j,\xi}(v),$$

where

$$q_{j,\xi}(\mathbf{v}) \coloneqq 2^{1/4} \psi_j\left(rac{\xi+2\mathbf{v}}{\sqrt{2}}
ight).$$

Vertical operators/poly-Fock space

Conclusions

 $\Omega = \mathbb{R}.$

For every
$$\xi$$
 in \mathbb{R} , $d_{\xi} = \dim(\widehat{H}_{\xi}) = m$.

For every
$$\xi$$
 in \mathbb{R} , $\left(q_{0,\xi}, \; q_{1,\xi}, \; \ldots, \; q_{m-1,\xi}
ight)$ is an orthonormal basis of $\widehat{H}_{\xi}.$

$$\mathcal{C}(\rho_{\mathcal{F}_m})\cong \mathcal{C}(\rho_{\mathcal{S}_m})\cong \mathcal{C}(\rho_{\mathcal{H}_m})=\mathcal{V}\cong L^\infty(\mathbb{R},\mathcal{M}_n(\mathbb{C}))\cong L^\infty(\mathbb{R})\otimes \mathcal{M}_n(\mathbb{C}).$$

Vertical operators in $\mathcal{F}_m \cong$ bounded matrix-functions on \mathbb{R} .

Possible themes for future works

- Vertical operators in $\mathcal{F}_m(\mathbb{C}^n)$.
- Angular operators in the wavelet spaces.
- More generally, shift-invariant operators associated to coherent states.
- Quasi-hyperbolic and quasi-nilpotent case in $\mathcal{A}^2(\mathbb{B}_n, \mu_{n,\alpha})$.
- Compute directly the Fourier transform of the reproducing kernel of the *m*-poly-Bergman space on the upper half-plane.
- Radial operators in many RKHS over the unit disk.